



TELGATE

A fungicide for the control of botrytis in grapes, boysenberries, strawberries, lemons, oranges and mandarins.

ACTIVE INGREDIENT: contains 500 g/litre fenhexamid in the form of a suspension concentrate (SC)





TELGATE®

HSNO CLASSIFICATION: Designed for biocidal action.

ENVIRONMENT: Designed for biocidal action. Do not apply into or onto water. Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area. Avoid contamination of any water supply with product or empty container. PERSONAL PROTECTION: When mixing or loading wear overalls,

impervious gloves and eye/ face protection. Avoid contact with or inhalation of soray mist. Wash hands and exposed skin after work.

STORAGE: Store in original container tightly closed and in a secure, dry, cool, well ventilated area, away from foodstuffs. Storage must be in accordance with the current version of NZS 8409 Management of Agrichemicals.

Shelf life: When stored appropriately this product should show no significant degradation for 2 years from the date of manufacture. Contact the supplier for further information about the use of this product if it is older than this.

SPILLS AND DISPOSAL: When dealing with spills wear personal protective clothing and equipment as described in the PERSONAL PROTECTION section. Contain spilled product and store in labelled, sealed drums for safe disposal. If spray mixture is spilled absorb with sawdust, soil or other suitable material.and collect in drums as above. Deal with all spillages immediately. If contamination of drains, streams, watercourses, etc. is unavoidable, warn the local water guthority.

PRODUCT DISPOSAL: If possible dispose of by using according to the label, otherwise dispose of in an approved landfill.

CONTAINER DISPOSAL: Triple rinse and add rinseate to the spray tank.
Take to an Agrecovery site or burn if circumstances, especially wind direction, permit otherwise burn in a suitable land fill.

FIRST AID:

If medical advice is needed, have product container or label at hand. For advice call a doctor or the National Poisons Centre (0800 764 766). IF SWALLOWED: Rinse mouth. Do NOT Induce vomiting. Call a POISON CENTRE or doctor for advice.

IF ON SKIN: Wash with plenty of soap and water.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If irritation persists, get medical advice.

IF INHALED: Remove to fresh air and keep at rest in a position comfortable for breathing.

SAFETY DATA SHEET: A safety data sheet is available from Grosafe Chemicals Ltd.

EMERGENCY RESPONSE: 0800 CHEMCALL (0800 243 622)

GENERAL INFORMATION:

TELGATE® is a contact fungicide for the control of botrylis in grapes, boysenberries, strawberries, lemons, oranges and mandarins. Fenhexamid prevents fungi from infecting plants by inhibiting spore germination and mycelial growth. The fungicide is absorbed into the outer waxy layer of plant surfaces and is protected from being washed off by rainfoll for irrigation.

RESISTANCE MANAGEMENT:

GROUP 17 FUNGICIDE

TELGATE® contains fenhexamid which belongs to the hydroxyanalide class of fungicides. Resistance to this fungicide could develop in some situations from repeated use. To minimise this risk, use strictly in accordance with label instructions and resistant management strategies that exist for any disease listed on the label. Do not use more than 2 applications of TELGATE® in the same crop during the same season.

TELGATE® should be used in a spray programme that incorporates fungicides with different modes of action. Refer to the NZCPR website (www.nzpps.org/resistance/index.php) for more detailed information.

DIRECTIONS FOR USE:

It is an offence to use this product on animals.

Note: Shake the container well before use. For larger containers roll or invert the container before use.

Mixing: Add the required amount of TELGATE® to a partly filled spray tank, then add the remainder of the water while thoroughly agitating. Continue agitation while spraying.

Compatibility: TELGATE® is compatible with most commonly used insecticides and fungicides.

RECOMMENDATIONS

The rates recommended are for high volume spraying to the point of run-off. For concentrate spraying adjust dilution rate accordingly.

CROP	DISEASE	RATE	REMARKS
Grapes	Botrytis	75ml/100 litres or a minimum of 750ml/ha	Make two applications at pre bunch closure and veraison, or at veraison and again 14 days later. Apply in sufficient water to achieve good coverage of the bunches.
Boysenberries Strawberries	Botrytis	150ml/100 litres or a minimum of 1.5 litres/ha	Make two applications over the flowering period ensuring good coverage of the flowers.
Lemons Oranges Mandarines	Botrytis	100ml/100 litres	Make up to two applications as a full cover spray to point of run off at full open flowering.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

WITHOLDING PERIODS:

Grapes – 21 days Boysenberries – 35 days Strawberries – 1 day Lemons and oranges – 7 days

Clementine/Satsuma Mandarins - Do not apply after the end of flowering

Registered pursuant to the ACVM Act 1997, No P9576 See www.foodsafety.govt.nz for registration conditions. Approved pursuant to HSNO Act 1996, Approval Code: HSR000008 See www.epa.govt.nz for approval conditions

Registered and distributed by:



20 Jean Batten Drive
 Mt Maunganui
 P.O. Box 14-450 Tauranga 3141
 Tel: 0800 220 002

Tel: 0800 220 002 Email: info@grosafe.co.nz www.arosafe.co.nz GSTG170/B02

WARRANTY: This formulation is warranted to contain the specified active ingredient within accepted analytical tolerance when packed. The user bears the risk for damage resulting from factors bevond the manufacturer's control. The manufacturer declines all liability for damage

resulting from improper storage or use of the product.

BATCH No: DATE OF MANUFACTURE:

TELGATE® is a Registered Trade Mark of Grosafe Chemicals Ltd